

# Stabilized R-Lipoic Acid



*The highly effective form of lipoic acid*

By Cristiana Paul, M.S.

THIS INFORMATION IS PROVIDED FOR THE USE OF PHYSICIANS AND OTHER LICENSED HEALTH CARE PRACTITIONERS ONLY. THIS INFORMATION IS INTENDED FOR PHYSICIANS AND OTHER LICENSED HEALTH CARE PROVIDERS TO USE AS A BASIS FOR DETERMINING WHETHER OR NOT TO RECOMMEND THESE PRODUCTS TO THEIR PATIENTS. THIS MEDICAL AND SCIENTIFIC INFORMATION IS NOT FOR USE BY CONSUMERS. THE DIETARY SUPPLEMENT PRODUCTS OFFERED BY DESIGNS FOR HEALTH ARE NOT INTENDED FOR USE BY CONSUMERS AS A MEANS TO CURE, TREAT, PREVENT, DIAGNOSE, OR MITIGATE ANY DISEASE OR OTHER MEDICAL CONDITION.

## Lipoic Acid Chemical Forms: the R and S Isomers

Alpha-Lipoic Acid (or thioctic acid) is synthesized from the amino acid cysteine and octanoic acid in plants and animals as both the R and S isomers, with a great majority as R-lipoic acid and only trace amounts of S-lipoic acid. When alpha-lipoic acid is synthesized in a laboratory, a 50/50 racemic mixture of the R and S enantiomers\* results in what we call R/S lipoic acid, which is the most popular form of commercially available lipoic acid (also named RAS lipoic acid or all racemic lipoic acid). Our patients know it as simply alpha lipoic acid. New technological advancements have allowed for the stabilization and isolation of the RLA (R-lipoic acid isomer), which is believed to account for all the benefits seen from studies with R/S lipoic acid. Supplementing with pure RLA, thus eliminating the presence of SLA (S-lipoic acid isomer), creates new opportunities for intensive nutritional support because SLA is believed to interfere with the effectiveness of RLA.<sup>16-21, 25</sup>

## How is Stabilized R-Lipoic Acid Different From Other R-Lipoic Products on the Market?

R-lipoic acid is a highly unstable compound that easily polymerizes into a sticky rubber or glue-like substance if it is not prepared, stored and processed correctly. This may adversely affect bioavailability. Stable RLA is a non-hygroscopic, non-polymeric potassium salt form of RLA. The capsules are heat stable, characterized by fast dissolution rates, high solubility and absorption. They are also free of residual solvents and moisture.

There are wide variations in the R/S ratios and total polymer contents of raw material and finished products now on the market. Few supplement companies have experience with RLA and are unaware of the myriad of problems associated with its encapsulation and stability. Designs for Health's **Stabilized R-Lipoic Acid** is one of the most stable and highly bioavailable forms of R-lipoic acid available on the market.

## R-Lipoic Benefits

Lipoic acid is a disulfide compound that is a cofactor in vital energy-producing reactions in the body. It is also a potent biological antioxidant, both water and fat soluble. It is made endogenously in humans and so it is not an essential nutrient. However, many physiological states, such as excessively high blood glucose levels, diabetic polyneuropathy, cataract, liver pathologies and toxic metal load, make lipoic acid conditionally essential. In addition, extensive research indicates that the many roles of alpha-lipoic acid may result in various health benefits, as reviewed here.

\*Enantiomers are a set of molecules identical in composition yet with a different spatial conformation which confers them different chemical and physiological properties.

R/S-lipoic acid has been studied for over 30 years and it is approved in Germany as a drug for the treatment of polyneuropathies, such as diabetic and alcoholic polyneuropathies, and liver disease.<sup>1</sup>

Recent studies have investigated the effectiveness of RLA versus SLA forms in order to identify their specific effects. It was suspected that RLA, being the majority of the natural form produced in the body, would have a stronger impact than SLA and the results of the research have confirmed this.<sup>16-21, 25</sup>

## Bioavailability

Compared to SLA, RLA causes 50% higher peak plasma levels of lipoic acid and 60-85% higher total absorption.<sup>12</sup> Feeding lipoic acid to animals at risk of cataract caused a 2-7 fold higher uptake of RLA versus SLA in the lens content of lipoic acid, and reduced the development of experimentally produced cataract by 50%.<sup>19</sup>

## Boosts Energy Production/Mitochondrial Cofactor

RLA is the majority of lipoic acid found in nature and therefore likely to fit better as a cofactor for mitochondrial enzymes pyruvate and alfa-ketoglutarate dehydrogenase.<sup>5</sup> SLA cannot bind well to these enzymes and actually inhibits them.<sup>20</sup> Thus the S-form can oppose the action of the R-form. In the aging rat heart, RLA stimulated ATP production, whereas SLA inhibited it.<sup>3</sup>

RLA supplementation improves metabolism, measured as oxygen consumption in liver cells, and improves ambulatory activity in supplemented animals, bringing old treated animals to the level of young animals.<sup>3</sup> Pre-treatment of brain cells with RLA leads to the restoration of the mitochondrial activity lost due to glutathione depletion.<sup>6</sup>

## Enhanced Glucose Metabolism

RLA significantly increases insulin sensitivity, glucose transport, metabolic rate and reduces the gain in body fat associated with aging.<sup>10-11</sup> R-lipoic acid has insulin-mimetic effects in glucose uptake in insulin resistant cells and may have therapeutic implications in restoring glucose availability in tissues such as the skeletal muscle.<sup>13, 16</sup>

## Supplement Facts

Serving Size 1 capsule

Amount Per Serving	% Daily Value	
Biotin	4 mg	1333%
Taurine	500 mg	*
R-Lipoic Acid	100 mg	*

\*Daily Value not established.

**Other Ingredients:** Potassium bicarbonate, microcrystalline cellulose, silicon dioxide, vegetable stearate.

The RLA was found to enhance insulin-stimulated glucose transport and non-oxidative/oxidative glucose metabolism by as much as 64%, while SLA had no effect. Also, RLA decreased insulin by 17% while SLA increased it by 15%.<sup>16</sup>

Through its positive effects on cellular energy metabolism, RLA attenuates metabolic dysfunction associated with advanced glycation end products (AGEs). AGEs accumulate on long-lived proteins, including beta-amyloid plaques in Alzheimer's disease and contribute to neuronal dysfunction and cell death.<sup>21</sup>

### Antioxidant Properties

RLA increases cellular and mitochondrial antioxidant activity, and was able to eradicate the age-related changes in animal models. This effectively attenuates the reported age-related increase in oxidative stress.<sup>3</sup>

RLA significantly increases or recycles other antioxidants including coenzyme Q10, vitamin C, vitamin E and glutathione.<sup>3, 5, 6, 11</sup> RLA protects lipids against peroxidation and reverses stress damage in the heart.<sup>7</sup>

### Anti-inflammatory Effect

RLA, a membrane permeable antioxidant, prevents the up-regulation of the AGE-induced gene expression responsible for regulating nitric oxide (NO) production. NO oxidizes nitrates and proteins which are markers of a chronic neuroinflammatory condition. This mechanism is relevant for Alzheimer's disease and for many chronic inflammatory conditions.<sup>24</sup> RLA reduces inflammation, and is more potent by a factor of 10 over R/S-LA.<sup>17</sup>

### Metal Chelator

RLA was more effective than the SLA in a battery of metal chelation tests. One hypothesis of the cause of diabetic complications involves overloading by transition metals, which implicates the RLA as more effective in treating diabetic neuropathy.<sup>2</sup>

### Neuroprotection

RLA improves memory, reverses cognitive dysfunction, and protects the brain from neurodegeneration associated with aging. This may be due to its effect on increased ATP production, chelating, antioxidant and anti-glycating capacity.<sup>6, 7</sup>

### Summary of SLA (S-Lipoic Acid) Properties

Until recently it was believed that SLA was physiologically inactive. Results from studies performed in vitro and with animals comparing the R with the S or R/S LA are warranting the use of pure RLA over the racemic ALA, whenever possible. SLA cannot bind with critical mitochondrial enzymes and inhibits ATP production.<sup>20</sup> At high concentrations, S-lipoic acid inhibits mitochondrial metabolism. It is metabolized in the outer cell membrane or cytoplasm which may interfere with RLA's ability to penetrate the inner mitochondrial membrane, thus limiting energy production.<sup>20</sup> SLA does not improve glucose disposal and slightly increases insulin levels.<sup>16</sup>

SLA has some positive benefits, but no advantage over the pure R form. SLA can function as an antioxidant but it recycles 38 times slower than RLA.<sup>14</sup> It also has some metal chelating properties, yet inferior to the R form.<sup>16</sup> Although toxicity studies have proven SLA to be safe up to very high levels, it is clear that SLA is not a metabolically preferred molecule.

### References

1. Alpha-lipoic acid as a biological antioxidant. Packer L, Witt EH, et al. *Free Rad Biol Med* 1995;19:227-250.
2. Thioctic (lipoic) acid: a therapeutic metal-chelating antioxidant? Ou P, Tritschler HJ, Wolff SP. *Biochem Pharmacol* 1995; 50:123-126.
3. (R)-alpha-Lipoic acid-supplemented Old Rats Have Improved Mitochondrial Function, Decreased Oxidative Damage, and Increased Metabolic Rate. Hagen TM, Ingersoll RT, Ames AB. *FASEB J* 1999 13:411-418.
4. Influence of selegiline and lipoic acid on the life expectancy of immunosuppressed mice. Freisleben HJ, Neeb A, Lehr F, Ackermann H. *Arzneimittelforschung* 1997 Jun; 47(6): 776-80.
5. Pharmaceutical composition containing R-alpha-lipoic acid or S-alpha-lipoic acid as active ingredient. Ulrich H, Weischer CH, et al. US Patent 5,728,735, 1998.
6. Pre-treatment with R-lipoic acid alleviates the effects of GSH depletion in PC12 cells: implications for Parkinson's disease therapy. Bharat S. *Neurotoxicology*. 2002 Oct; 23 (4-5):479-86.
7. Oxidative stress in the aging rat heart is reversed by dietary supplementation with (R)-alpha-lipoic acid. Suh JH, Shigeno ET, et al. *FASEB J* 2001 Mar; 15(3): 700-6
8. Memory loss in old rats is associated with brain mitochondrial decay and RNA/DNA oxidation: partial reversal by feeding acetyl-L-carnitine and/or R-alpha-lipoic acid. Liu J, Head E, et al. *Proc Natl Acad Sci U S A* 2002 Feb 19; 99(4): 2356-61.
10. Oral administration of RAC-alpha-lipoic acid modulates insulin sensitivity in patients with type-2 diabetes mellitus: a placebo-controlled pilot trial. Jacob S, Ruus P, et al. *Free Rad Biol Med* 1999 Aug;27 (3-4):309-14.
11. R-alpha-Lipoic Acid Action on Cell Redox Status, the Insulin Receptor, and Glucose Uptake in 3T3-L1 Adipocytes. Moines H, Trios O, et al. *Arch Biochem Biophys* 2002 Jan 15; 397(2): 384-91.
12. Breithaupt-Grogler K, Niebh G, et al. Dose-proportionality of oral thioctic acid--coincidence of assessments via pooled plasma and individual data. *Eur J Pharm Sci*. 1999 Apr;8(1):57-65.
13. Cytokine-induced glucose uptake in skeletal muscle: redox regulation and the role of alpha-lipoic acid. Khanna S, Roy S, Packer L, Sen CK. *Am J Physiol*. 1999 May;276(5 Pt 2):R1327-33.
14. Biewenga GP, Dorstijn MA, et al. Reduction of lipoic acid by lipoamide dehydrogenase. *Biochem Pharmacol*. 1996 Feb 9;51(3):233-8.
15. Age-associated decline in ascorbic acid concentration, recycling, and biosynthesis in rat hepatocytes--reversal with (R)-alpha-lipoic acid supplementation. Lykkesfeldt J, Ames BN, et al. *FASEB J* 1998 Sep; 12(12): 1183-9.
16. Differential effects of lipoic acid stereoisomers on glucose metabolism in insulin-resistant skeletal muscle. Streeper RS, Henriksen EJ, et al. Tritschler HJ. *Am J Physiol* 1997 Jul; 273(1 Pt 1): E185-91.
17. Ulrich H, Weischer CH, et al. Pharmaceutical composition containing R-alpha-lipoic acid or S-alpha-lipoic acid as active ingredient. US Patent 5, 728, 735, 1998.
18. Cytosolic and mitochondrial systems for NADH- and NADPH-dependent reduction of alpha-lipoic acid Haramaki N, Han D, et al. *Free Radic Biol Med* 1997; 22(3): 535-42
19. Stereospecific effects of R-lipoic acid on buthionine sulfoximine-induced cataract formation in newborn rats. Maitra I, Serbinova E, et al. *Biochem Biophys Res Commun* 1996; 221:422-429.
20. Stereospecific effects of lipoic acids on mammalian pyruvate dehydrogenase complex. Hong YS, Jacobia SJ, Packer L. *Free Radic Biol Med*. 1999 Mar;26(5-6):685-94.
21. Neuroprotection by the metabolic Antioxidant alpha Lipoic Acid. Packer, L; Tritschler, H; Wessel K, 1997 *Free Rad Biol Med* 22, Nos 1/2, 359-378.
22. Mitochondrial decay in the aging rat heart: evidence for improvement by dietary supplementation with acetyl-L-carnitine and/or lipoic acid. Hagen TM, Moreau R, Suh JH, Visioli F. *Ann N Y Acad Sci*. 2002 Apr;959:491-507.
23. Advanced glycation endproducts induce changes in glucose consumption, lactate production, and ATP levels in SH-SY5Y neuroblastoma cells by a redox-sensitive mechanism. de Arriba SG, Loske C, et al. *Cereb Blood Flow Metab*. 2003 Nov;23(11):1307-13.
24. Anti-inflammatory antioxidants attenuate the expression of inducible nitric oxide synthase mediated by advanced glycation endproducts in murine microglia. Wong A, Dukic-Stefanovic S, et al. *Eur J Neurosci*. 2001 Dec;14(12):1961-7.
25. Interaction of alpha-lipoic acid enantiomers and homologues with the enzyme components of the mammalian pyruvate dehydrogenase complex. Loeffelhardt S, Bonaventura C, et al. *Biochem Pharmacol*. 1995 Aug 25;50(5):637-46.