


BioPC Pro



CLINICAL APPLICATIONS

- Supports Healthy Cellular Structure, Function and Signaling
- Supports Mitochondrial and Immune Function
- Supports GI Barrier Health and Integrity
- Helps Maintain Normal Inflammatory Balance

IMMUNE HEALTH[†]

Phospholipids are an integral part of biological membranes, and they tend to decrease as we age. They are critical for optimal cell and mitochondrial membrane function, including their growth, shape and repair. Optimal cell membrane function facilitates healthy cell signaling, which is important for energy production, immune function and regulation of inflammatory cascades. Additionally, BioPC Pro leverages the benefits of a full-spectrum phosphatide blend to help balance the nervous system and bolster parasympathetic tone in patients who may be sympathetic dominant. Together, phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylinositol (PI) and phosphatidylserine (PS) help achieve optimal nervous system function when it is overstimulated. BioPC Pro also helps digest and process fats and supports better focus, cognitive function, and overall brain health. This rapidly absorbed blend provides 5,900 mg* of naturally bioavailable phosphatides per serving, making it one of the most concentrated preparations available.

*Subject to natural variability

Overview

Phosphatides make up the major structural lipids in cellular membranes.¹ PC was one of the first biological amphiphiles (a molecule having both hydrophilic and hydrophobic parts) to be discovered, and it is the most abundant phospholipid in cellular membranes.^{2,3} With this initial research conducted at the same time the fluid mosaic cell surface model was developed, it led to a widespread view of phosphatides as cellular building blocks.¹

But emerging evidence is making the case that, in addition to its structural role, phosphatides also play a significant biological

role in cell signaling, maintaining mitochondrial function, and transforming cellular membranes, enzymes, and receptors to be more functional.^{1,4-10} While phosphatides are sometimes used individually, the use of more complex phosphatide blends containing PC, PE, PI, and PS are considered more useful.^{1,11-13} Research shows that oral phospholipids are processed efficiently by the body, with more than 90% absorbed within hours of ingestion.^{1,14,15} For these reasons and more, restoring phosphatide levels through supplementation can be highly beneficial.

Phosphatides and the Mitochondrial-Immune Interface

In addition to cellular energy production, mitochondria perform several critical functions in the cell that make phospholipid movement to and from the mitochondrion essential for cellular integrity, including the modulation of calcium signaling, regulation of cell death, and maintenance of cellular redox balance and innate immune signaling.^{1,2,16}

While phosphatides make up a large portion of the lipids comprising cellular and intracellular membranes, it is their metabolism that creates a tightly controlled cell signaling network essential for a healthy innate immune response.¹⁷ Phosphatides and their metabolites influence multiple aspects of innate immunity, including cell shape, stickiness and degranulation.³ Phospholipids themselves are also recognized by innate-like T cells, which are considered essential for recognizing foreign organisms, as well as self-antigens.³

When various cellular stressors, including reactive oxygen species (ROS) and danger signals, are released in the presence of cellular or mitochondrial damage, the formation

[†] These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



of inflammasomes are induced, which triggers innate immune defenses.^{1,18-20} Inflammasomes such as NLRP3 induce cytokine release, which increases oxidative stress and inhibits autophagy and mitophagy.^{1,21,22} The NLRP3 inflammasome is, itself, activated by mitochondrial ROS, suggesting that healthy mitochondria are essential for a balanced immune response.¹

Studies show that stealth immune challenges are also associated with elevated mitochondrial oxidative stress.²³ These results suggest an interruption in cellular communication may be at the core of continued mitochondrial stress and danger signals in stealth immune challenges.²³ Oral phosphatides can be used to help modify mitochondrial signals and promote healthy inner mitochondrial membrane potential, thereby improving electron transport chain function and ATP synthesis (including perceived energy, mood and cognition) and overall immune-health balance.^{1,24,25}

Phosphatides and the Gut-Immune Interface

In a healthy gastrointestinal tract, the presence of PC in the mucosal secretion mucus creates a hydrophobic surface layer that safeguards the intestinal tissues and modulates mucosal signaling.^{1,26,27} PC represents more than 90% of the phospholipids comprising intestinal mucus, which serves as a primary component of the intestinal barrier, creating a protectant shield against GI-related immune challenges by binding foreign matter.²⁸ Poor nutrition, stress insults and alterations in the gut microbiome all have a negative impact on the health of intestinal mucus.²⁹ In the presence of GI challenges, the PC content of intestinal mucus may decrease by up to 70%, leaving the distal GI tract especially vulnerable. However, several studies show that oral PC supplementation replenishes the phosphatide content of intestinal mucus, reinforcing the body's first, and perhaps most important, immune defense mechanism.³⁻⁵

Directions

1 scoop (10.2 grams) per day with a meal or as recommended by your health care professional. May be mixed in juice or shakes, or sprinkled on food.

Does Not Contain

Gluten, corn, yeast, artificial colors or flavors.

Cautions

If you are pregnant or nursing, consult your health care professional before taking this product.

Supplement Facts ^{v1}		
Serving Size 1 Scoop (10.2 Grams)		
Servings Per Container About 30		
	Amount Per Serving	% Daily Value
Calories	60	
Total Fat	5 g	6%*
Saturated Fat	1 g	5%*
Polyunsaturated Fat	3 g	
Monounsaturated Fat	1 g	
Total Carbohydrate	3 g	1%*
Protein	<1 g	
Phosphorus	130 mg	10%
Sunflower Lecithin Powder	10 g	**

* Percent Daily Values are based on a 2,000 calorie diet.
** Daily Value not established.

Other Ingredients: Silicon Dioxide and Mixed Tocopherols.

ID# 173030 10.8 oz (306 Grams)

Typical naturally occurring phospholipid profile (per 10 g serving)†:

Phosphatidylcholine 2.5 g (2,500 mg)
 Phosphatidylinositol 2 g (2,000 mg)
 Phosphatidylethanolamine 1 g (1,000 mg)
 Phosphatidic acid 0.4 g (400 mg)
 †subject to natural variability

References

- Nicolson GL, Ash ME. Lipid Replacement Therapy: a natural medicine approach to replacing damaged lipids in cellular membranes and organelles and restoring function. *Biochim Biophys Acta*. 2014;1838(6):1657-1679.
- Furse S, de Kroon AI. Phosphatidylcholine's functions beyond that of a membrane brick. *Mol Membr Biol*. 2015;32(4):117-119.
- Acoba MG, Senoo N, Claypool SM. Phospholipid ebb and flow makes mitochondria go. *J Cell Biol*. 2020;219(8):e202003131.
- Blaton V, Vandamme D, Peeters H. Activation of lipoprotein lipase in vitro by unsaturated phospholipids. *FEBS Lett*. 1974;44(2):185-188.
- Horsch AK, Hudson K, Day AJ. Uptake and metabolism of 3H-fatty acid labelled lecithin by normal and atherosclerotic intima in vivo and in vitro. *Atherosclerosis*. 1977;26(4):493-504.
- Howard AN, Patelski J. Hydrolysis and synthesis of aortic cholesterol esters in atherosclerotic baboons. Effect of polyunsaturated phosphatidyl choline on enzyme activities. *Atherosclerosis*. 1974;20(2):225-232.

† These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

EFFICACY
the power of *e*

7. Waligora Z, Patelski J, Brown BD, Howard AN. Effect of a hypercholesterolaemic diet and a single injection of polyunsaturated phosphatidyl choline solution on the activities of lipolytic enzymes, acyl-CoA synthetase and acyl-CoA cholesterol acyl-transferase in rabbit tissues. *Biochem Pharmacol.* 1975;24(24):2263-2267.
8. Karaman A, Demirbilek S, Sezgin N, Gürbüz N, Gürses I. Protective effect of polyunsaturated phosphatidylcholine on liver damage induced by biliary obstruction in rats. *J Pediatr Surg.* 2003;38(9):1341-1347.
9. Olbrich K, Rawicz W, Needham D, Evans E. Water permeability and mechanical strength of polyunsaturated lipid bilayers. *Biophys J.* 2000;79(1):321-327.
10. Buko V, Artsukevich A, Maltsev A, et al. Effect of polyunsaturated phosphatidylcholine on lipid structure and cAMP-dependent signal transduction in the liver of rats chronically intoxicated with ethanol. *Exp Toxicol Pathol.* 1994;46(4-5):375-382.
11. Nicolson G. Lipid replacement as an adjunct to therapy for chronic fatigue, anti-aging and restoration of mitochondrial function. *JANA.* 2003;6(3):4-10.
12. Nicolson GL, Ellithorpe R. Lipid replacement and antioxidant nutritional therapy for restoring mitochondrial function and reducing fatigue in chronic fatigue syndrome and other fatiguing illnesses. *Journal of Chronic Fatigue Syndrome.* 2006; 13(1):57-68.
13. Nicolson GL. Lipid replacement/antioxidant therapy as an adjunct supplement to reduce the adverse effects of cancer therapy and restore mitochondrial function. *Pathol Oncol Res.* 2005;11(3):139-144.
14. Zierenberg O, Grundy SM. Intestinal absorption of polyenephosphatidylcholine in man. *J Lipid Res.* 1982;23(8):1136-1142.
15. Zierenberg O. Clinical and biochemical studies of the transport of polyenephosphatidylcholine in human serum and its physiological impact on cholesterol distribution between serum lipoproteins. *Phospholipids and Atherosclerosis.* Avogaro P (ed). Raven Press, New York. 1983:175-189.
16. Cloonan SM, Choi AM. Mitochondria: sensors and mediators of innate immune receptor signaling. *Curr Opin Microbiol.* 2013;16(3):327-338.
17. O'Donnell VB, Rossjohn J, Wakelam MJ. Phospholipid signaling in innate immune cells. *J Clin Invest.* 2018;128(7):2670-2679.
18. Brookes PS, Yoon Y, Robotham JL, Anders MW, Sheu SS. Calcium, ATP, and ROS: a mitochondrial love-hate triangle. *Am J Physiol Cell Physiol.* 2004;287(4):C817-C833.
19. Latz E, Xiao TS, Stutz A. Activation and regulation of the inflammasomes. *Nat Rev Immunol.* 2013;13(6):397-411.
20. Dostert C, Pétrilli V, Van Bruggen R, Steele C, Mossman BT, Tschopp J. Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science.* 2008;320(5876):674-677.
21. Franceschi C, Bonafè M, Valensin S, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci.* 2000;908:244-254.
22. Rodgers MA, Bowman JW, Liang Q, Jung JU. Regulation where autophagy intersects the inflammasome. *Antioxid Redox Signal.* 2014;20(3):495-506.
23. Peacock BN, Gherezghiher TB, Hilario JD, Kellermann GH. New insights into Lyme disease. *Redox Biol.* 2015;5:66-70.
24. Agadjanyan M, Vasilevko V, Ghochikyan A, et al. Nutritional supplement (NT Factor™) restores mitochondrial function and reduces moderately severe fatigue in aged subjects. *Journal of Chronic Fatigue Syndrome.* 2003;11(3):23-36.
25. Seidman MD, Khan MJ, Tang WX, Quirk WS. Influence of lecithin on mitochondrial DNA and age-related hearing loss. *Otolaryngol Head Neck Surg.* 2002;127(3):138-144.
26. Dial EJ, Zayat M, Lopez-Storey M, Tran D, Lichtenberger L. Oral phosphatidylcholine preserves the gastrointestinal mucosal barrier during LPS-induced inflammation. *Shock.* 2008;30(6):729-733.
27. Eehalt R, Braun A, Karner M, Füllekrug J, Stremmel W. Phosphatidylcholine as a constituent in the colonic mucosal barrier--physiological and clinical relevance. *Biochim Biophys Acta.* 2010;1801(9):983-993.
28. Stremmel W, Eehalt R, Staffer S, et al. Mucosal protection by phosphatidylcholine. *Dig Dis.* 2012;30 Suppl 3:85-91.
29. Diebel LN, Liberati DM. Reinforcement of the intestinal mucus layer protects against *Clostridium difficile* intestinal injury in vitro. *J Am Coll Surg.* 2014;219(3):460-468.